

*Thesis for Doctor of Philosophy Degree in the Faculty  
of medicine*

*Bundelkhand, University,  
Jhansi (India)*

**A** *study of childhood Epilepsy and Epileptic Syndromes in a  
subset of population in India with emphasis on hereditary  
factors.*

*Dr Brahm Prakash*

*Supervisor  
Dr N.S.Sengar  
M.D., D.M.*




*Institution: Bundelkhand university Jhansi (India)*

*Place Of study: Northern Railway Central Hospital,  
New Delhi  
&  
Bundelkhand, Jhansi (India)*


*Thesis*

*Entitled*

***"A study of childhood Epilepsy and Epileptic Syndromes in a subset of population in India with emphasis on hereditary factors"***



***Dr Brahm Prakash***  
***Candidate***



***Dr N.S. Sengar***  
***M.D., D.M.***  
***M.L.B. Medical College, Jhansi***  
***Supervisor***


***Place of study: Northern Railway Central Hospital, New Delhi***  
***&***  
***Bundelkhand University,***  
***Jhansi (India)***

*Thesis  
Entitled*


**A** *study of childhood Epilepsy and Epileptic Syndromes in a subset of  
population in India with emphasis on hereditary factors.*

*Was submitted to the Faculty of medicine, University of Bundelkhand  
University, for the degree of Doctor of Philosophy*

*On .....*



*Dr Brahm Prakash  
Candidate*



*Dr N.S. Sengar  
M.D., D.M.  
Supervisor*

*Bundelkhand University, Jhansi (India)*

***Declaration***

***I declare this study is substantially my own original work and has not been submitted in any form for an award at any other academic institution.***

Signature.....

Dr Brahm Prakash



## *List of Contents*

<b>1. Introduction and Review of literature.....</b>	<b>1</b>
Seizure in pediatric age group.....	3
Classification of Seizures.....	3
The Syndromes.....	5
Incidence.....	7
Prevalence.....	8
Febrile convulsions.....	9
Hereditary factors in Epilepsy.....	11
 <b>2. Aims and Objectives.....</b>	 <b>19</b>
 <b>3. Material and Methods.....</b>	 <b>21</b>
Geographical area of study.....	23
 <b>4. Data and Observations.....</b>	 <b>25</b>
Male -Female Ratio.....	26
Classification Seizure Type.....	28
Anti-Epileptic Medicines.....	30
Computer Tomography.....	32
Magnetic Resonance Imaging.....	34
Percentage of Pediatric Epilepsy cases.....	36
Agewise distribution of Seizures cases in children.....	38
Syndromic Classification of Seizure cases as per ILAE-89.....	40
 <b>5. Discussion.....</b>	 <b>44</b>
 <b>6. conclusion.....</b>	 <b>50</b>
 <b>7. Recommendation.....</b>	 <b>53</b>
 <b>8. Referances.....</b>	 <b>55</b>
 <b>9. Appendix.....</b>	 <b>64</b>
Performa used for the study of epilepsy in the children.....	65
Seizure Classification used in the study ILAE-89.....	66
Map of Railway network showing geographical area of probands.....	67

## **Acknowledgement**

*I would like to avail this opportunity to place on record my heart felt thanks to all those who helped me in completion of this study.*

*I express my sincere gratitude to the authorities of Bundelkhand University and Prof. Ramesh Chandra in particular in allowing me to pursue this study in children.*

*I am extremely grateful to Dr N.S. Sengar for guiding and supervising the work without whose suggestions and overall supervision it was not possible to complete this study.*

*I am grateful to Dr K.K. Gupta Medical director NRCH for continuous support and encouragement for the study of epilepsy in children at central hospital New Delhi.*

*I am thankful to Dr D.K. Das CMD/NR for providing excellent facilities for Anti-Epileptic drug serum levels and neuroimaging studies without which it was not possible to conduct this study.*

*I do not have words and means to repay the debt of Prof. Satish Jain in introducing me to the research methodology in epilepsy during my earlier work "Classification of Seizures in children".*

*I will be failing in my duty if I do not mention the encouragement and support which I got from Dr K.Suresh D.G./RHS for my study. It was his persistent motivation, which encouraged me to take up this work.*

*I do not have words to express my thanks to my sons Vaibhav and Anubhav. They have never complained in my devoting the time for this study, which I should have given to them. I am also thankful to them for resquing me whenever I had problem on computer while compiling this work.*

*Last but not the least it was not possible to complete this study without constant love, care and strength which my wife Vinod gave me. In fact she was the force behind motivating me constantly to compete this work.*

*I am thankful to Dr Charan Singh, Shri M S Rana and Shri Ashok Kumar Bhardwaj helping me in completion of this work.*

*Dr Brahm Prakash*

**Introduction & Review of Literature**

*A study of childhood Epilepsy and Epileptic Syndromes in a subset of population in India with emphasis on hereditary factors.*

## **II. Generalized seizures**

- *Absence seizures (typical or atypical)*
- *Myoclonic seizures*
- *Clonic, tonic or tonic-clonic seizures*
- *Atonic seizures.*

## **III Unclassified seizures.**

- *Whether partial or generalised*

The limitation of this classification was it was confined to an individual seizure type. Seizure is an event with which a patient comes to a physician but the condition to which this seizure belongs is the epileptic syndrome and it is also the language in which two physicians can communicate. The International Classification of epileptic seizures and epileptic syndromes 1985/1989 (Commission on Classification and Terminology of the International League against epilepsy. Proposal for revised classification of epilepsies and epileptic syndromes. *Epilepsia* 1989,30; 389-399) tried to overcome the shortcomings of the previous classifications.

This Classification was exciting because in recognizing electroclinical syndromes early in the diagnosis of epilepsy, it was possible to predict response to treatment and more so the prognosis.

### ***The syndromes:***

*Syndromes are clusters of signs and symptoms customarily occurring together and include:*

- *The clinical picture of seizure*
- *The age of patient at the time first seizure*
- *EEG*
- *Evolution and prognosis*
- *Associated neurological features*
- *Positive or negative family history*
- *Neuroimaging investigative studies.*

*Two dichotomies are used in this classification-*

*Localization related onset or generalised*

*Etiology known or unknown:*

*Idiopathic: no underlying causes, may be inherited*

- *Symptomatic:*  
*As a consequence of known or a suspected disorder of CNS*
- *Cryptogenic:*

*Cause is hidden or occults whereas in there do not fulfil the criteria for idiopathic while on the other hand there is no proof of their symptomatology.*

*The symptomatic epilepsies constitute about 48% while 52% are idiopathic in nature. The incidence of symptomatic is higher then reported in the developed countries (Oka E, Ishida S, Ohtsuka Y, Othara S. Classification of epilepsies and epileptic syndromes of childhood according to the 1989 ILAE classification. J epilepsy 1993; 6:272-6). This is mainly because of higher incidence of perinatal insult and CNS infections. The insults to the nervous system in the neonatal period account for nearly one third of all symptomatic cases (Eriksson KJ, Koivikko MJ. Prevalence, Classification, and severity of epilepsy and epileptic syndromes in children. Epilepsia 1997; 38 : 1275-82)*

*The other cause of symptomatic epilepsy are tumors, intracranial granulomas, trauma, vascular strokes, AV malformation, congenital malformation, inborn error of metabolism etc (Viani F, Beghi E, Atza G, Gulotta MP. Classification of epileptic syndromes: advantage and limitations for evaluation of childhood epileptic syndromes in clinical practice. Epilepsia 1988; 29:440-5.)*

*Over the last decade, epidemiological studies have demonstrated that the prognosis of many childhood epilepsies is more favorable with 70-80 % reaching remission. However sever cases, some of which are refractory to treatment account for 20-30% of all cases (Eriksson KJ, K Koivikko MJ. Prevalance, classification and severity of epilepsy and epileptic syndromes in children. Epilepsia 1997; 38:1275-82).*

## ***Incidence:***

*Incidence is the rate at which new cases of a condition occur in a population. The incidence of epilepsy for all age group is about 40/100,000 persons years (Hauser WA, Angers JF, kurlandLT. The incidence of epilepsy in Rochester, Minnesota 1935-79. Epilepsia 1984;25:666).*

*The age specific incidence of unprovoked seizers in virtually all studies is highest in the first month of life. Seizure incidence then remains relatively high but it declines during the first year of life, and tends to decrease slightly thereafter through the*

second decade, after which incidence remains constant at about 30/100,000 in the adult years, only to increase again after the age of 60.

Contrary to common opinion, only about 50% of all cases of epilepsy start in childhood. In most studies, about 60% to 70% of cases are of unknown cause and in childhood, the majority of new cases in children involve generalized seizures from onset. Most, total population studies of incidence report a slight male preponderance.

In a study by W. Allen Hauser from National Institute of Neurological Disorders and Stroke, Bethesda, the large pediatric population was followed from the first prenatal visit of the mothers in pregnancy until children born of those pregnancies were 7 years old. The study was sampled by the Collaborative Perinatal Project (NCPP) of the National Institute of Neurological and Communicative Disorders and Stroke, provided valuable information regarding the epidemiology of epilepsy. In that population, 8.0 white children per thousand had at least one nonfebrile seizure, not considered being symptomatic of acute neurologic illness, between the ages of 1 month and 7 years. This is similar to the cumulative incidence for one or more unprovoked seizures of 1.1% by age 10 in the population of Rochester Minnesota. In black children in the NCPP study, the cumulative incidence rate was 9.0 per thousand. The difference between the racial groups was not statistically significant. Seizures occurred with similar frequency in boys and girls of the two racial groups. Minor motor seizures and neonatal seizures were not different in frequency by race or sex. (Clinical aspect of pediatric epilepsy volume 56 suppl., part 2)

Age-specific incidence rate (Rochester, Minnesota, 1935-1979)

Age group	No patients	rate/100,000	proportion	Male/Female
Years		Persons-years	Idiopathic%--Partial%	%
<1	41	121	54	17
1-4	85	63	81	21
5-14	129	44	84	47
				120
				97
				123



## **Prevalence:**

Prevalence of epilepsy is a measure of the proportion of patients currently suffering from active epilepsy or experiencing the consequences of epilepsy as measured by taking anticonvulsant. Since one need to identify existing cases of the illness, prevalence studies are considerably easier to conduct than incidence itself. The estimated prevalence of epilepsy varies widely in reported studies by a factor of 20, and ranges from a low of 2/ 100 0 in Marianas islands (Stanhope JM, Brody JA, Brink E. Convulsions among the Chamorro people of Guam, Marianas Islands. I. Seizure disorders *Am J Epidemiology* 1972; **95**: 292-298) to a high of 37/10,000 in Nigeria (Osunto kun Bo, Aevja AOJ, Nottidge VA, et al. Prevalence of the epilepsies in Nigerian Africans: a community-based study, *Epilepsia* 1987; **28**:272-279). Unfortunately this variation seems more related to study methodology and to definitions of epilepsy in individual studies than to true population variations. In prevalence studies based upon incidence cohorts, prevalence tends to increase with advancing age through early childhood, only to stabilize in the teen age and young adult years (Hauser and Nelson, *Epidemiology of epilepsy*; Cleveland clinic journal of Medicine: Volume 56 Suppl. Part 2).

## **Febrile convulsions:**

Febrile convulsions are the commonest type of genetic abnormality. 5% of all children will have a fit under five years of age, 3% of whom fulfil the criteria for febrile convulsions. The incidence is higher in Japan at 7-8% (Forfar and Arneil's Textbook of Paediatrics 1998 fifth edition, Churchill Livingstone, London).

Febrile convulsions rarely develop in to epilepsy, and they spontaneously remit without specific therapy. They have uniformly excellent prognosis. Febrile seizures are age dependent and are rare before 9 months and after 5 years of age. The peak age of

onset is 14-18 months of age, and incidence approaches 3-4% of young children. A strong family history of febrile convulsion in sibling and parents suggests a genetic predisposition. Linkage studies in several large families have mapped the febrile seizure gene to chromosome 19p and 8p 13-21. An autosomal dominant inheritance pattern is demonstrated in some families (Nelson Textbook of Pediatrics 16th edition, Harcourt Asia pte Ltd. 2000, Singapore.)

**Risk of recurrence:** 33% of all children with febrile convulsion will have recurrence of these 9% will have >3 recurrences, 75% of recurrences would occur within 1 year of first episode and 90% within 2 years (Nelson KB, Ellensburg JH. Prognosis in children with febrile seizures. Pediatrics 1978; 61:702-7.) The best predictor is early age of seizure and positive family history of febrile seizure (Berg AT, Shinar S, Hauser WA, ET AL. Predictors of recurrent febrile seizure: a meta-analytic review. Pediatric 1990; 116: 329-37).

### **Risk of subsequent epilepsy:**

The overall risk of epilepsy after febrile seizure is 2-2.5% (Verity CM, Golding J. Risk of epilepsy after febrile convulsion: A national cohort study. Br Ed J 1991; 303: 1373-6.) The prognosis is more guarded if the convulsion is prolonged or atypical. Up to 40% may have another convulsion and 15% a third episode. If the child suffers from multiple repeated febrile convulsions the possibility of early malignant epilepsy such as myoclonic epilepsy of Dravet exists.

The risk factors for subsequent epilepsy are

- The child is less than 12 months old;
- Complex febrile seizure occurred with neurological sign
- Prolonged seizure lasts more than 30 minutes.
- Developmental/ neurological abnormality present before first seizure.
- There are more than three episodes
- Non febrile seizure in parents/ siblings.

convulsions. This was the case in an analysis of the Rochester, Minnesota, data set (Rich et al., 1987)

There may well be a subset of children who have an autosomal -dominant mode of inheritance of febrile seizures (Joshin et al.,) and Rich et al., 1987. At this time no definitive identification of a gene or locus for febrile seizure has been established. Genes on chromosomes 8 (Wallace et. al., 1996) and 19 (Johnson) have been linked to some cases of febrile seizures in large families. Rapid advances in this area in future are anticipated. Most likely all children have some increased susceptibility to seizures from fever at the specific age window. Genetic influences are therefore likely to account for some but not all of the cases.

There is little doubt that epileptic seizures tend to aggregate in families (Hauser WA et al- clinical Aspects of Pediatric Epilepsy -Cleveland Clinic Journal of Medicine Vol. 56 Suppl. Part 2pp115-193). While a number of diseases follow Mendelian pattern of inheritance and have as part of their manifestation the occurrence of seizures, these conditions, in aggregate, will account for no more than 1% of seizures in childhood (Anderson VE, Hauser WA, Rich SS, Genetic heterogeneity in the epilepsies (In Degado-Escueta AV, Ward AA JR, Woodbury DM, Porter RJ, eds The mechanism of epilepsy. New York, Raven Press, 1986,pp59-75.). Risk for epilepsy is increased by a factor of three for individuals with a first degree relative with epilepsy, an overall risk similar to that associated with head injury or infection of the central nervous system (CNS)(Annegers J F, Hauser WA, Anderson VE, kurland LT, The risk of seizure disorders among relatives of patients and childhood onset epilepsy Neurology 1982; 32; 174-179). Similarly, risk for epilepsy is increased by a factor of three for children with a sibling who has had febrile seizure (Annergers JF, Hauser WA, Anderson VE, Kurland LT. The risk of seizure disorders among relatives of patients with childhood onset epilepsy, Neurology 1982; 32:174-179). While there is a perception that generalized - onset seizures are associated with higher risk for epilepsy in relatives, this seems a result of very high risk among siblings with epilepsy manifested by absence seizures or Myoclonic seizures. If these unique but rare subgroups are excluded, risk for epilepsy among relatives of probands with epilepsy characterized by generalized - onset seizures are similar to those in relatives of probands with epilepsy manifested by partial

seizures. Family history is important in modifying risk for epilepsy even in the presence of a history of overt cerebral insult.

In the study by Collaborative Perinatal Project (NCP) of the National Institute of Neurological and communicative Disorders and stroke, familial factors were also predictors of seizure disorder in children. Family history factors showing significant univariate association with seizure disorders included maternal seizure disorders, maternal MR, Paternal congenital malformations, and seizures or motor deficits in older siblings. Paternal seizures were not observed to be predictive; however, histories were taken from mothers who might not always have been acquainted with the fathers medical histories except in one mother child pair with tuberous sclerosis, specific heritable disorders were not recognized (Nelson KB, Ellen Berg JH. Predisposing and causative factors in childhood epilepsy. *Epilepsia* 1987; 28(suppl); S16-S24).

The familial tendency is thought to be carried on chromosome 6 and to be age dependent in its manifestation. Unless EEG studies with hyperventilation and photic provocation are carried out between 3 and 15 years one may miss the dominant transmission, as clinical seizure may not occur. Precipitation of seizure by computer game has reinforced this as an important area. Lennox 50 years ago showed that the number of the close relatives with an epileptic EEG was six times higher with known epileptics than with normal controls (Forfar and arneil's Text book of Pediatrics, Churchill Living stone 5th edition pp677-680).

Genetic diseases with seizures a symptom there are more than 200 Mendelian conditions with epilepsy as main symptom. Tuberous sclerosis is a good example of how a genetic disease may present as any one of the three clinical form i.e. genetic, lesional or malignant.

- There is incomplete penetrance so uncomplicated epilepsy may occur in members of the family without any other clinical manifestation of the diseases.
- Single or multiple small gliomas, hamartomas and tubers act as focal 'lesions'.

- Malignant epilepsy such as west's syndromes or Lennox-Gastaut syndromes can occur in the infant of a tuberous sclerosis mother, especially after several generations. Dominant inheritance causes worse disease when acquired from the mother and may result in brain malformation with cortical dysphasia.

Some EEG studies suggest that the generalized spikes and wave EEG pattern in the inherited as an autosomal dominant trait with age-dependent expression. Other workers have suggested that polygenic inheritance is a better explanation of the data. The risk for sibling of patient with generalized epilepsy ranges from 4% to 8% which is substantially lower than would be expected from a simple gene disorder. Partial epilepsies have often been regarded as nongenetic or lesional types of epilepsy. However, certain types do show clear genetic influence. These include benign partial epilepsies of childhood and posttraumatic epilepsy.

Sibling and offspring of the affected probands have three to four fold increase in the febrile convulsion rate compared with the general population. Interpretation of the family studies on febrile convulsions has resulted in the conflicting conclusions. The inheritance said to be autosomal dominant, autosomal recessive or polygenic. The relationship of the febrile convulsion to epilepsy is complex and there is some evidence that there may be two genetically distinct subtypes of febrile convulsions with different risk of the subsequent epilepsy. In addition many children with febrile convulsion will subsequently show generalized spikes and wave on EEG even though they do not have convulsion, as do a high proportion of siblings.

By discovering a linkage between juvenile myoclonic epilepsy and HLA markers, the gene responsible was successfully mapped to human chromosome 6p (Durner et al 1991). The locus was subsequently designated EJM1. than by virtue of the close association of juvenile myoclonic epilepsy and other generalized epilepsy disorder (childhood pyknolaptic absence epilepsy (typical absence epilepsy), grand mall Seizures on early morning awakening, photo convulsive television or video games epilepsy), together with the high incidence of EEG abnormality in the close relatives, investigators concluded that the gene imparts a tendency for abnormal electrical activity to synchronize and generalize in histologically normal cerebral cortices.



The various epileptic syndromes are actually the different ways in which this tendency expresses itself (Delgado- Escueta et al 1990, Greenberg et al 1995). Juvenile myoclonic epilepsy can be considered the prototype, which has a mixture of all three types of absence, myoclonic and generalized tonic-clonic fits, while the rest are milder variants. The expression of the gene is highly age dependent, just, as childhood absence epilepsy tends to resolve after adolescence. EEG can not reliably do family screening. As it may be absent in older individuals linkage or DNA analysis may be necessary.

A recent report of a linkage study showed no evidence for a locus on chromosome 6p in some British and Swedish families with juvenile myoclonic epilepsy and primary Grandmall seizures (Whithouse et al 1993). It is therefore reasonable to conclude that there is more than one gene carrying the tendency for generalized fits. It is interesting to note that valproic acid is an effective anti epileptic drug for these syndromes. It is different from drugs in that it is a fatty acid with eight carbon molecules. Eight carbon fatty acids are anticonvulsant and ten carbon fatty acids are anaesthetic agents. Octanoic acid was thought to be the fatty acid with anti epileptic properties in the ketogenic diet.

The fact that valproic acid appears specific for many genetic epilepsies suggest that it may correct an underlying metabolic abnormality. Hopefully gene cloning will provide insight in to the mechanism of the fits as well as the anti epileptic action of valproic acid. (Forfar and Arneil's Text book of Pediatrics, fifth edition, Churchill living stone, pp681 to 682

Hypoglycemia, hypocalcaemia or hypomagnesemia will not be included in this group. These are acute dysrhythmias, which complicate acute encephalopathy, which may be recurrent, and not simply the seizure. Conditions such as mitochondrial encephalopathy with lactic acidosis and stroke like episodes (MELAS syndrome). And Myoclonic epilepsy with ragged red fibers (MERRF) is included. These disorders have either a known enzyme defect or morphological evidence of storage abnormality but the enzyme defect is still unknown. They are almost invariably associated with other neurological abnormality, particularly intellectual deterioration. Thus the presence of family history with a symptomatic combination of seizure and mental retardation should

trigger a search for inborn errors of metabolism as this has implications for treatment and genetic counseling.

Following is the list of genetic epilepsies:

- **-Benign familial neonatal seizure – AD:**  
Chromosome 20, nonsense mutation of acetylene receptor  
(leppert et al 1989)  
Normal development
- **-Idiopathic partial epilepsy of infancy-50% family history:**  
related to dominant neonatal fit  
Neurology normal  
Others on chromosome 19
- **-Benign myoclonic epilepsy of infancy**  
-AR seen in identical twins,  
valproate sensitive,  
normal development.
- **-Childhood Absence epilepsies -AD:**  
variable penetrance and age  
nearly normal Dependent  
expression; 60-70% female
- **-Photosensitive epilepsy:**
  - Genetically inherited and
  - Amenable to valproate
  - Normal neurology
  - More common in females
  - Photosensitivity persists inspite of treatment
- **-Epilepsy with myoclonic absence**
  - Male preponderance
  - Neurology exam normal,
  - mental Retardation in 45 to 75% cases

- *Juvenile absence epilepsy*
  - Good response to Valproate*
  - Normal neurology*
- *Juvenile myoclonic epilepsy*
  - Chromosome 6,*
  - Excellent response to Valproate,*
  - Normal Neurology*
- *Benign rolandic epilepsy :*
  - with variable penetrance age dependent*
  - Absence of neurological and intellectual deficit*
- *Benign occipital with occipital spike-waves*
  - Difficult to treat*

*Source: Disorders of central nervous System, Forfar and Arneil's Textbook of Pediatrics Churchill Living stone fifth edition 1998, pp680-681.*

*A study by S.Jain, MV Padma, A Puri, Jyoti and MC Maheshwari from the department of Neurology, Neuroscience's center All India Institute of Medical sciences New Delhi India.*

*Large numbers of families with many members having seizures have been used to understand the role of hereditary factors in the pathogenesis of human epileptic syndromes. The aim was to establish genetic database to form a hypothesis on the possible genetic contributions in different epileptic syndromes. The study concluded that A significant percentage (19%) of first and second degree relatives of probands with all types of epileptic syndromes have seizures*

- *The risk of relatives being affected varied as a function of relation with the proband. Concordance of epileptic syndromes between probasnds and relatives was related to the epileptic syndromes in probands.*



- The syndrome of SSEL is probably a benign epileptic syndrome seen in Indians genetically predisposed to seizures. Hereditary factors may play an almost equal role in the predisposition of relatives of epilepsy in families of probands with different epileptic syndromes.

*In another study by S. Jain, Menka S. Jain, M V Padma, A Puri, P. Sen and M C Maheswari from department of Neurology, Neuroscience's Center AIIMS New-Delhi India on Epilepsies among twins born in families of Indian probands with epilepsy had following conclusions-*

1. The-twinning rate among families of Indian probands with epilepsy is similar to that seen in data from hospital births in the same city and other studies reported from India

2. Epilepsy in twins may be largely genetic rather than due to factors associated with twinning

3. Family data such as in this study, if collected meticulously, can be used to form hypothesis for understanding the extent of contribution by genetic factors towards the pathogenesis of complex genetic diseases like human epilepsies.

*In another study by S. Jain, Padma M V, Tripathi M, Narula A, Seema, Gujreet and MC Maheswari on Phenotypic analysis of JME:*

Implication for gene discovery strategies the conclusion was that persons diagnosed with JME having absence (7%) overlap with absence epilepsy. JME affected having PPR on EEG (8%) and those needing VPA and another drug for seizure control (10%) could be the other subtypes. Among all JME there could be about 25% who may have not the classical syndrome. The familial and sporadic cases could be other subgroups. These possible sub types could be responsible for confounding result of

*genetic studies (Paper presented at the Asia and Oceanian Congress 11-13Nov. 2000  
New Delhi India)*

***Aims and objectives of the study***

- 1. To classify the seizure types in the children among the Family members of railway employees*
- 2. To know the incidence of past history of febrile convulsions in the Children presenting with different seizure types*
- 3. To know the occurrence of seizures among family members.*

## ***Material and methods***


*The present study was carried in the department of Pediatrics at Northern Railway Central Hospital, New Delhi. The entire study group consisted of Railway Health Services beneficiaries. Two hundred and eighty six children who attended NRCH, out patient department or indoor department or admitted through the causality in the age group up to 16 years were included in the study.*

*The Northern Railway Central hospital is referral hospital for whole of Northern Railway and surrounding area of other Zonal Railways. The map on the next page shows the geographical area from where these patients have been coming to central hospital.*

*The geographical areas of Punjab, Haryana, Utter Pradesh and some parts of Rajasthan and Delhi as a whole are included in Northern Zone.*

*The map showing geographical area of residence of patients is shown on the next page:*

# RAILWAY NETWORK OF INDIA



Map not to Scale  
Copyright (c) Mapience India Limited 2001

*The Railway workforce is permanent employees of the organization are stable population as continuation of the job up to sixty years of age make a more suitable group for prolonged follow up of the probands and their families. The follow up is easier than city mobile population. The railways give comprehensive treatment to all Railway employees. The transportation up to and for attending NRCH is free. All investigations including neuroimaging studies and serum level studies for AED levels and pharmaceuticals are free irrespective of the cost.*

*Railway population is subset of population in India where the socioeconomic status, permanent employment total free and readily accessible quality comprehensive health care is available to all the beneficiaries, There is no waiting for specialist advice at NRCH. All patients requiring indoor admission or OPD consultation are given the same on same day. Or the availability of health facilities is available to them at par of any developed country in the world*

*All patients were attended between september 1998 to 1999 were included in the study. Complete clinical details were recorded. History of febrile seizure was included in clinical details of the patients. The details were recorded on a Performa.*

*Performa has been annexed.*

*Details of EEG were recorded. EEG facility is available in NRCH. All EEG's were personally reported by trained Neuro-Psychiatrist of the hospital.*

*The classification of epilepsies and epileptic syndromes were done on the basis of revised classification given by International commission on classification and terminology of the International League Against Epilepsy 1989. The classification of ILAE-89 is annexed.*

*The family pedigree was constructed to include all first degree and second-degree relatives of the probands. Seizures in the family members were documented and effort was made to examine all the available affected relatives.*

*The seizure classification and entire family data was verified on every visit of the patient.*



- ***Data and observations***

**Male- Female ratio**


*Total two hundred and eighty six cases have been included in the study.*

*One hundred and sixty eight probands belonged to male sex.*

<i>Total cases</i>	<i>286</i>
--------------------	------------


<i>Female patients</i>	<i>118</i>
------------------------	------------

<i>Male patients</i>	<i>168</i>
----------------------	------------




**Classification: Seizure Type****History based :**

1.	<i>Focal/partial /sec generalized Sz.</i>	<i>096</i>
2.	<i>Generalized tonic or clonic or tonic clonic Sz.</i>	<i>142</i>
3.	<i>Juvenile Myoclonic epilepsy</i>	<i>010</i>
4,	<i>Childhood absence epilepsy</i>	<i>020</i>
5.	<i>West syndrome</i>	<i>004</i>
6.	<i>Single Sz</i>	<i>005</i>
7.	<i>Febrile seizures</i>	<i>009</i>



<i>Anti Epileptics used</i>	
<i>Sodium Valproate</i>	<i>034</i>
<i>Clobazam</i>	<i>009</i>
<i>DPH</i>	<i>023</i>
<i>Carbamezepine</i>	<i>200</i>
<i>Combination</i>	<i>020</i>






<u><b>Computed Tomography done</b></u>	
<i>Total CT done</i>	<i>168</i>
<i>Normal</i>	<i>58</i>
<i>Abnormal</i>	<i>110</i>




### Computed Tomography of cranium with contrast



<i>Magnetic Resonance (MRI)</i>	
<i>Total MRI</i>	<i>10</i>
<i>Normal</i>	<i>03</i>
<i>Abnormal</i>	<i>07</i>


Magnetic Resonance



Series1	10	3	7
---------	----	---	---


<i>Percentage of epilepsy cases in total pediatric</i>	
<i>Patients attended at NRCH</i>	
<i>Total pediatric patients attended NRCH during this period-</i>	<i>6234</i>
<i>Total number of seizure cases in children</i>	<i>286</i>
<i>Percentage of epilepsy cases among total pediatric</i>	
<i>Patients attending NRCH</i>	<i>4.57%</i>

## percentage of Epilepsy cases in Paerdiatric patients attending NRCH



<u><i>Age wise distribution of seizure disorder in children</i></u>	
<i>0-5 years</i>	<i>72</i>
<i>6-10 years</i>	<i>131</i>
<i>11-16 years</i>	<i>83</i>
<i>Total children attended at NRCH for seizure disorder</i>	<i>286</i>






Age wise distribution of seizure disorder in children

0-5 years	72
6-10 years	131
11-16 years	83

<u>Syndromic classification of seizures after EEG, Nueroimaging studies and other required investigation (Classification as per ILAE-89)</u>	
1. Localization related- epilepsies and syndromes	107
2. Generalized epilepsies and syndromes	120
3. Epilepsies and syndromes undetermined,	017
Weather localized or generalized Focal/ Generalized, Neonatal Sz	
4. Special syndromes. Febrile Sz, Single Sz, Situation related Sz.	0 12
5. Seizure with single Small Enhancing CT Lesion (SSEL)	030





Syndromic classification of seizures after EEG, Neuroimaging and other investigations in protocol  
Classification as per ILAE-89

1. Localisation related epilepsies and syndromes	107
2. Generalised epilepsies and syndromes	120
3. Epilepsies and syndromes undetermined	17
4. Special syndromes-Single Sz, Febrile Sz, Situational Seizures	12
5. Seizures with single Small ring Enhancing CT lesion	30

<i>Percentage of febrile seizures</i>	
<i>Total no of cases of epilepsy</i>	<i>286</i>
<i>History of febrile seizures present in</i>	<i>15</i>
<i>%age of febrile seizures in the study group</i>	<i>5.24</i>

<i>Cases where family history was positive in the family of Probands</i>	
<i>Total cases of Epilepsy</i>	<i>286</i>
<i>Family history positive in</i>	<i>60</i>

**Discussion**

*I*t has been estimated that 50 percent of epileptic cases has its onset in pediatric age group. In our study out of total six thousand and thirty children who attended the pediatric dept of NRCH from Sept.1998 to Aug 1999. Incidence of epilepsy for all age group is about 40/100,000 person years (Hauser WA, Annegers JF, Kurland LT. The incidence of epilepsy in Rochester, Minnesota 1935-79, *Epilepsia* 1984; 25:666). There were two hundred and eighty six cases of epilepsy, which are approximately 4.67 % of total attendance NRCH in Pediatric department. It is less in comparison to developed countries because our beneficiaries are habituated in a linear fashion along the Railway track and big proportion do not come to NRCH for obstetric care during child birth. This the main reason that in our study the maximum cases of epilepsy are in age group of 6 to 10 years age bracket. In all other studies the percentage of first Sz is maximum in 0 to 1 year (Peter R. Camfield and Carol S. Camfield *Pediatric epilepsy: an over view* chapter 37: pp637).

In our study the number of male affected children is higher which is compatible to the existing literature.

Seizure types have been classified on the basis of history and clinical examination first and followed by using Syndromic classification given by International League Against Epilepsy in 1989(*Epilepsia* 1989) after work up of cases like compulsory EEG, Haemogram, CXR, CT/MRI if required and fundus examination (in every case), Mauntonox test and Serum IgG etc for Neurocysticercosis if required. Serum levels were done for levels of anti epileptic medicines where seizure control was poor. Sodium Valproate has been used where Idiopathic generalized epilepsy was diagnosed as absence or Myoclonic type and where carbamazepine was not leading to good seizure control. The Carbamazepine has been main stay of treatment in most general tonic clonic type of seizure disorder and partial seizures. Cost of medicine and availability was not the issue as Railways free of cost supply all pharmaceuticals. The follow up was quit regular because the transport and medicine and all investigations to railway beneficiaries are free. The availability of specialist for consultation is easily accessible on all working days of the hospital. The admission to children is not denied even if the wards are over crowded. The specialist service to the patients is available to patients on all days and odd hours also through emergency department of the hospital.

The syndromic classification has been useful for managing the epileptic cases for following reasons

- Judicious uses of anti epileptic drugs like Sodium Valproate in absence and Myoclonic Epilepsy.
- When to start AED.
- When to withhold the medicine.
- When to withdraw medication: for example in Benign Centro-temporal epilepsy one can withdraw the medicine early or even withhold the AED but one has to continue medicine for prolonged period in Occipital epilepsy.
- To predict the prognosis for example in Juvenile myoclonic epilepsy AED has to be for prolonged or life long need. The prognosis of West Syndrome is bad in comparison of Benign Rolandic epilepsy.

The Syndromic classification gives more accuracy in diagnosis like EEG in Absence epilepsy where 3 Hz per sec spike is almost the signature of the disease with this one can be accurately administer AED (Sodium Valproate) and give prognosis of the disease. If EEG slow spike wave paroxysms of less than 2.5 Hz on an abnormal background activity, commonly is associated with mental retardation as is seen in Lannox Gastot Syndrome. ILAE classification of 1989 which appeared in *Epilepsia* 1989; 30:389-399 is syndromic classification. The syndromes are clusters of signs and symptoms customarily occurring together and include-

- Clinical event
- Age of onset
- Ictal and interictal EEG
- Evolution and prognosis
- Associated neurological features
- Family history
- Neuroimaging

In our studies idiopathic generalized epilepsies are more as in other studies of India and abroad one is of Oka ET AL (*J Epilepsy* 1993; 33:1072-7). The symptomatic epilepsies are more in India because of incidence of perinatal insult and CNS infections and SSEL. Probands had a family history comparable to that those of GES. The syndrome of SSEL



*appears to be benign epileptic syndrome seen in the Indian population that is genetically predisposed to seizures. (S.Jain ET alEpilepsia.Vol.38.no, 1997)*

*The use of ILAE classification 89 was used in our study in two hundred and eighty six cases presented to us in NRCH. The classification was useful in management and predicting the prognosis of the cases. It is recommended that every children pediatric neurology division should adopt in protocol to classify the seizures, This will improve the quality of management of epilepsy in children. There is some need to make classification more users friendly as this classification is complex and many syndromes are not adequately defined. The other limitations of seizure classification is that it describes common seizure phenomenology only and can not reliably distinguish partial from generalized origin of seizures is a crude guide to selection of drug treatments as per Dr Engels "A major contribution of the International League against Epilepsy was the establishment of standardized classification and terminology for epileptic seizures and syndromes. This provides a universal vocabulary that not only facilitated communication among clinicians, but also established a taxonomic foundation for the performance of quantitative clinical and basic research on epilepsy. The Executive committee of the ILAE, which took office in July in 1997, is reviewing the current classification we now await a new user-friendly classification, which may include pathophysiological Substrates, underlying gene defects and functional disability due to epilepsy.*

### ***Febrile seizures:***

*In our study nine cases were purely of febrile seizures, that is approximately three percent of the study group. The children with the epileptic syndromes who had history of febrile convulsions were only 1% in our study group. These children were separate then purely febrile seizure children who were in age group of 3 years to 5 years of age proves that febrile seizure is not a major risk factor for subsequent epilepsies in children. In our study only ten MRI were done so no comment can be given regarding association between Mesial Sclerosis and febrile convulsions. However studies shows that (Camfield PR, Camfield CS. Management of febrile seizures. Current problems in Pediatrics 1997; 27(1): 6-13) Mesial temporal sclerosis is only 1/75000 children. In our study group febrile patients were managed by intermittent prophylaxis with oral diazepam .It seems that counseling to alley the anxiety of parents should be main for this benign disorder.*



### ***Family history of seizure disorder in the family of probands:***

*In our study group of epileptic children sixty cases were having positive family history in first degree or second-degree relatives of the probands. This is 60% of the study group, which is comparable to other studied in this geographical area. Family history was more positive in the children who were having idiopathic generalized seizure disorder. The family history was also more common in the children who had EEG finding of Centrotemporal spikes (Benign Centro-temporal epilepsy)*

*The children who had febrile convulsions also had family history positive proving febrile seizure disorder a genetic inherited benign disorder. In one study at AIIMS by Satish Jain et al the family history in the first degree and second degree relatives was found to be 19% (S. Jain et al Occurrence of Epilepsies in family members of Indian probands with different epileptic syndromes; Epilepsia: 38(2): 237-244.1997).*

*In our study the family history is positive in sixty cases, which is approximately 20.97% of the study group.*

## **Conclusion**

*In this study we have included two hundred and eighty cases of epilepsy that were presented to us in the department of Pediatrics at NRCH New Delhi from September 1999 to August 2000. The children were presented to us through casualty, Pediatric OPD or were admitted to the indoor department after referral from peripheral referral centers of Indian Railways. The geographical area from where children came has been Delhi and surrounding areas of Delhi, Railway population residing in Punjab, Uttar Pradesh, Haryana, Rajasthan and Utteranchal states of India. The study used ILAE-1989 classification and found that it is useful for management of the patients for following reasons*

- *It allows the judicious use of ant-epileptic drugs (AED)*
  - When to start AED*
  - When to withdraw AED*
  - And help to decide when to withhold the medicine*
- *In predicting the prognosis of epilepsy in a particular child by knowing his classified group as per ILAE-89.*
- *For defining the likelihood of identifying the underlying pathology.*

*In this study fifteen cases of febrile seizures were found in the total study group. The percentage of febrile seizures is same as found in general population in other studies. We conclude that febrile seizure is not a very significant factor in leading to epilepsy in children in later life. The family history in the cases of febrile seizures was present in significant numbers indicating it to be a genetic inherent disorder.*

*The family history was meticulously recorded in all the cases. The pedigree was made up to second degree relatives of the Probands. It has been found that family history was positive in sixty children who presented with epilepsy to this referral center. This figure comes to 20.97% of cases, which is consistent with other studies in this part of the world. The study is indicative that in epileptic syndromes in children hereditary plays a very important role. In the present study the symptomatic cases of epilepsy are in significant numbers. It is because the perinatal insults and CNS infections like tuberculosis and protozoa infestations like Taenia Solium are quite prevalent in this part of the world.*

*In age wise distribution in total study group of two hundred and eighty six epileptic cases 72 were in 0-5 years, 131 were in 6-11 years and 83 were in 11-16 years of age group.*

*The number of children in 0-5 years is less in comparison to available data in other national and international studies. This is because our population is residing in a linear fashion in the cities, towns and stations along the Railway track of Indian railways. The seizures in children are maximum in first twenty eight days of life the neonates are managed locally as few mothers comes for deliveries at NRCH from distant cities and towns and prefer confinement near their normal place of residence. In higher age group children are brought by the parents when seizures are recurrent needs neuroimaging studies and other lab investigation like serum levels etc. This is a selection bias in our study because of different geographical distribution of our beneficiaries.*

**Recommendations**

- *The present study was done in a subset of population in India. The study group was small the further study in a larger set up is required.*
- *The study proves that Classification of seizures using ILAE-89 should be adopted by all pediatric Neurology clinics as it helps in appropriate choice of AED.*
- *Family pedigree up to second degree relative should be drawn in all seizure cases.*
- *Neuroimaging studies like CT/MRI should be part of protocol for all patients of epilepsy.*
- *The investigations to rule out CNS infections like Tuberculosis and protozoal infestations like Taenia Solium should be done.*
- *The febrile seizure is not a significant risk factor for epilepsy in later years of life.*
- *Family pedigree is helpful in pointing the underlying pathology of the diseases. The positive history also gives clue for choosing the AED.*

**REFERENCES**



1. Dreifuss FE. The epilepsies: clinical implication of the international classification. *Epilepsia* 1990; 31 (Suppl.3): S3-10.
2. Commission on Classification and Terminology of the international League Against Epilepsy. Proposal for revised clinical and electroencephalographic classification of epileptic seizures. *Epilepsia* 1981; 22:489-501.
3. Commission on Classification and Terminology of the International League Against Epilepsy. Proposal for revised classification of epilepsies and epileptic syndromes. *Epilepsia* 1989; 30: 389-399.
4. Shah KN, Rajadhyaksha SB, Shah VS, Shah NS, Desai VG. Experience with the international League Against Epilepsy. Proposal for revised classification of epileptic seizures (1981) and epilepsies and epileptic syndromes (1989) in epileptic children in a developing country. *Epilepsia* 1992; 33: 1072-7.
5. Oka E, Ishida S, Ohtsuka Y, Ohatahara S. Classification of epilepsies and epileptic syndromes of childhood according to the 1989 ILAE classification. *J.Epilepsy* 1993; 6:272-6.
6. Viani F, Beghi E, Atza G, Gulotta MP. Classification of Epileptic syndromes: advantage and limitations of evaluation of childhood epileptic syndromes in clinical practice. *Epilepsia* 1988; 29:440-5.
7. Verity CM, Ross EM, Golding J. Epilepsy in the first 10 years of life: findings of the child health and education study. *Br.Med J* 1992; 305: 857-61.
8. Eriksson KJ, Koivikko MJ, Prevalence, Classification, and severity of epilepsy and epileptic syndromes in children. *Epilepsia* 1997; 38:1275-82.
9. Roger J, Beareau M, Dravet CH, Dreifuss FE, Perret A, Wolf P, eds. *Epileptic syndromes in infancy, childhood and adolescence*. 2<sup>nd</sup> ed. London: John Libbey, 1992.
10. Engel J. Classification of the International League Against Epilepsy: Time for reappraisal. *Epilepsia* 1998; 39:1014-17.
11. Luders H, Acharya J, Baumgartner, et al. Semiological seizure classification. *Epilepsia* 1998; 39: 1006-13.
12. Febrile seizures: Long term management of children with fever associated seizures. Summary of a National Institute of Health Consensus Statement. *Br Med. J* 1980; 281:277-9
13. Nelson KB, Ellenberg JH. Prognosis in children with febrile seizures. *Pediatrics* 1978; 61:702-7.
14. Berg AT, Shinnar S, Shapiro ED, et al. Risk factors for a first febrile seizure: a matched case-control study. *Epilepsia* 1995;36(4): 334-41.
15. Rantala H, Uhari M, Hietala J. Factors triggering the first febrile seizure. *Acta Pediatric* 1995; 84(4): 407-10.
16. Bethune P, Gordon KG, Dooley JM, et al. Which child will have a febrile seizure? *Am J Dis Child* 1993; 47:35-9.
17. Nelson KB, Hirtz DG. Febrile seizures. In: *Pediatric Neurology. Principles and Practice*, Ed.Kenneth F. Swaiman. Bosby, Missouri 2<sup>nd</sup> ed. Vol.1, 1994 pp 565-70.
18. Vandenberg BJ. Studies on convulsive disorders in young children. IV. Incidence of convulsions among siblings. *Dev Med Child Neurol* 1974; 16:457-64.

19. Ellenberg JH, Nelson K. Febrile seizures and later intellectual performance. *Arch Neurol* 1978;35:17-21.
20. Berg AT, Shinnar S, Hauser WA, et al. Predictors of recurrent febrile seizures: a meta-analytic review. *Pediatrics* 1990; 116:329-37.
21. Offringa M, Derksen Lubsen G, Bossuyt PM, et al. Seizure recurrence after a first febrile seizure: a multivariate approach. *Dev Med Child Neurol* 1992; 34:15-24.
22. El Radhi AS, Banajet S. Effect of fever on recurrence rate of febrile convulsions. *Arch Dis Child* 1989; 64:869-70.
23. Berg AT, Shinnar S, Darefsky AS, et al. Predictors of recurrent febrile seizures. A prospective cohort study. *Arch Pediatric Adolescence Med*. 1997; 151(4): 371-378.
24. Berg At, Shinnar S, Hauser WA, et al. A prospective study of recurrent febrile seizures. *N Eng. J Med*. 1992; 327:1122-7.
25. Verity CM, Butler NR, Golding J. Febrile convulsions in a national cohort follow up from birth. Prevalence and recurrence in the first five years of life. *Br Med. J* 1985; 290:1307-10.
26. Verity CM, Golding J. Risk of epilepsy after febrile convulsions: A national cohort study. *Br Med J* 1991; 303:1373-6.
27. Camfield PR, Camfield CS. Management and treatment of febrile seizures. *Current Problems in Pediatrics* 1997; 27(1):6-13.
28. Watanabe K, Takahashi I, Negoro T, et al. Benign epilepsy of children with complex partial seizures following febrile convulsions. *Seizure* 1993, 2: 57-62.
29. Liu Z, Mikati M, Holmes GL. Mesial temporal sclerosis: pathogenesis and significance. *Pediatric Neurol* 1995; 12(1): 5-16.
30. Maher J, McLachlan RS. Is seizure duration the most important predictor of temporal lobe epilepsy? *Brain* 1995; 118:1521-1528.
31. Lorber J, Sunderland R. Lumbar Puncture in Children with convulsions associated with fever. *Lancet* 1980; 1: 785-86.
32. Annegers JF, Hauser WA, Shirts SB, et al. Factors prognostic of unprovoked seizures after febrile convulsions. *N Engl J Med* 1987; 316:494-8.
33. Sofijanov N, Emoto S, Kuturee M, et al. Febrile seizures: Clinical characteristics and initial EEG. *Epilepsia* 1992; 33:52-7.
34. Practice parameter- The neurodiagnostic evaluation of the child with a first simple febrile seizure. American Academy of Pediatrics. Provisional Committee on Quality improvement, Sub-Committee on Febrile Seizures. *Pediatrics* 1996; 7(5): 769-72.
35. Sheila Wallace. Febrile Seizures. In: *Epilepsy in children*. Ed. Sheila Wallace. Pub. Chapman and Hall Medical, London 1996, 185-198.
36. Morales A, Bass N, Lakesmith K. Computerized tomography and febrile seizures (Abstract) *Ann Neurol* 1992; 32:432.
37. Berg AT, Shinnar S. Complex febrile seizures. *Epilepsia* 1996; 37(2): 126-33.
38. Rutter N, Smales OR. Role of routine investigations in children with their first febrile convulsions. *Arch Dis Child* 1977; 52: 188-91.
39. Knudsen FU. Rectal administration of diazepam in solution in the acute treatment of convulsions infants and children. *Arch Dis Child* 1979; 54: 855-7.

40. Uhari M, Rantala H, Vainionpaa, Kurttila R. Effect of acetaminophen and low intermittent doses of diazepam on prevention of recurrences of febrile seizures. *J. Pediatr* 1995; 126:991-5.
41. Guidelines for the management of convulsions with fever. Joint Working Group of the Research Unit of the Royal College of Physicians and the British Pediatric Association *Br Med J* 1991; 303:634-6.
42. Hirata Y, Mizuno Y, Nakano M, et al. Epilepsy following febrile convulsions. *Brain Dev* 1985; 7: 75.
43. Takizawa K, Sumi K. Prophylactic therapy for febrile convulsions – recurrence of febrile convulsions and occurrence of subsequent non-febrile convulsions on continuous prophylactic medication. *Brain Dev* 1984, 6: 69.
44. Ellenberg JH, Nelson K. Febrile seizures and later intellectual performance. *Arch Neurol* 1978; 35: 17-21.
45. Wolf SM, Carr A, Davis DC, et al. The value of phenobarbital in the child who has had a single febrile seizure. A controlled prospective study. *Pediatrics* 1977; 59:378-85.
46. Faero O, Kastrup KW, Nelson EL, et al. Successful prophylaxis of febrile convulsions with phenobarbital. *Epilepsia* 1972; 13: 279-85.
47. Newton RW. Randomized controlled trial s of phenobarbitone and valproate in febrile convulsions. *Arch Dis Child* 1988; 63: 1189-91.
48. Herranz JL, Armijo JA, Arteaga R. Effectiveness and toxicity of phenobarbital, primidone and sodium valproate in the prevention of febrile convulsions controlled by plasma levels. *Epilepsia* 1984; 25:89-95.
49. Mamelie N, Mamelie JC, Plasse JC, et al. Prevention of recurrent febrile convulsions – a randomized therapeutic assay: sodium valproate, phenobarbital and placebo. *Neuropediatrics* 1984; 15:37-42.
50. Knudsen FU. Effective short term diazepam prophylaxis in febrile convulsions. *J. Pediatr* 1985; 106:487-90.
51. Rosman NP, Colton T, Labazzo J, et al. A controlled trial of diazepam administered during febrile illness to prevent recurrent of febrile seizures. *N Eng J Med* 1993; 329:79-85.
52. Uhari M, Rantala H, Vainiopara, Kurttila R. Effect of acetaminophen and low intermittent doses of diazepam on prevention of recurrence of febrile seizures. *J Paediatr* 1995; 126:991-5.
53. Auteret E, Billard C, Bertrand P, et al. Double blind randomized trial of diazepam Vs placebo for prevention of recurrence of febrile seizures. *J Pediatr* 1990; 117:490-5/
54. Knudsen FU. Febrile seizures – treatment and outcome. *Brain Dev* 1996; 18(6): 438-49.
55. Baumer JH, David TJ, Valentine SJ, et al. Many parents think their child is dying when having a first febrile convulsion. *Dev Med Child Neurol* 1981; 23: 462-4.
56. Schmitt BD. Fever phobia. *Am J Dis Child* 1980; 134: 176-81.
57. Tondi M, Carboni F, Deriu A, Manca S, Mastropaolo C. Intermittent therapy with clobazam for simple febrile seizures. *Dev Med Child Neurol* 1987; 29: 830-31.

58. Manreza ML, Gherpelli JL, Machado-Haertel LR, Pedreira CC, Heise CO, Diamant A, et al. Treatment of febrile seizures with intermittent clobazam. *Arq Neuropsiquiatr* 1997; 55(4): 757-61.
59. Annegers JF, Hauser WA, Coan SP, et al: A population based study of seizures after traumatic brain injuries. *N. Engl. J Med.* 338:20, 1998.
59. Baram TZ: Pathophysiology of massive infantile spasms: Perspective on the putative role of the brain adrenal axis. *Ann Neurol* 33:231, 1993.
60. Baram TZ, Mitchell WG, Tournay A, et al: High dose corticotropin (ACTH) versus prednisone for infantile spasms: a prospective, randomized blinded study. *Pediatrics* 97: 375, 1996.
61. Berg AT, Shinnar S, Darefsky AS, et al: Predictors of recurrent febrile seizures *Arch Paediatr Adolesc Med* 151:371, 1997.
62. Berkovic SF, Andermann F, Carpenter S, et al: Progressive myoclonus epilepsies: Specific causes and diagnosis. *N Engl J Med* 315:296, 1986.
63. Cross JH, Jackson GD, Neville BGR, et al: Early detection of abnormalities in partial epilepsy using magnetic resonance. *Arch Dis Child* 69:104, 1993.
64. Delgado -Escueta AV, Enrile-Bacsal FE: Juvenile myoclonic epilepsy of Janz. *Neurology.* 34: 285, 1984.
65. Devinsky O: Cognitive and behavioral effects of antiepileptic drugs. *Epilepsia* 36 (suppl 2):S46, 1995.
66. Duchowny M, Harvey AS: Pediatric epilepsy syndromes: An update and critical review. *Epilepsia* 37 (Suppl 1): S26, 1996.
67. Duchowny M, Harvey AS: Pediatric epilepsy syndromes: An update and critical review. *Epilepsia* 37 (suppl 1):S26, 1996
68. Duchowny MS: Complex partial seizures of infancy. *Arch Neurol* 44:911, 1987.
69. Gross-Tsur V, Manor O, van der Meore J: Epilepsy and attention deficit hyperactivity disorder: Is methylphenidate safe and effective? *Pediatr* 130:40, 1987
70. Hauser WA, Rich SS, Lee JRJ, et al: Risk of recurrent seizures after two unprovoked seizures. *N Engl J Med* 338:429, 1998.
71. Holmes GL: Partial seizures in children. *Pediatrics* 77:725, 1986.
72. Lagae LG, Silbertsein J, Gillis PL, et al: Successful use of intravenous immunoglobins in Landau-Kleffner syndrome. *Pediatr Neurol* 18:165, 1998.
73. Leach JP, Brodie MJ: Tiagabine. *Lancet* 351:203, 1998
74. Maher J, MacLachlan RS: Febrile convulsions in selected large families. A single major-locus mode of inheritance? *Dev Med Child Neurol* 39:79, 1997.
75. Minassian BA, Lee JR, Henbrick JA, et al: Mutations in a gene encoding a novel protein tyrosine phosphate cause progressive myoclonus epilepsy. *Nat Genet* 20:171, 1998.
76. Neville BGR: Epilepsy in childhood. *Br Med J* 315:924, 1997.
77. Offringa M, Bossuyt PMM, Lubsen J. et al: Risk factors for seizure recurrence in children from febrile seizures. A pooled analysis of individual patient data from five studies. *J. Pediatric* 124:574, 1994.
78. Sanmdramouli S, Robinson R, Tsaloumas M, et al: Retinal hemorrhages and convulsions. *Arch Dis Child* 76:449, 1997.

79. Singh NA, Charlier C, Stauffer D, et al: A novel potassium channel gene, KCNO2, is mutated in an inherited epilepsy of newborns. *Nat Genet* 18:25, 1998.
80. Verity CM: Do seizures damage the brain? The epidemiological evidence. *Arch Dis Child* 78:78, 1998.
81. Vinning EPG, Freeman JM, Pillas DJ, et al: Why would you remove half a brain? The outcome of 58 children after hemispherectomy – the Johns Hopkins experience 1968 to 1996. *Pediatrics* 100:163, 1997.
82. Wallace SJ: First tonic-clonic seizures in childhood. *Lancet* 349:1009, 1997. Epilepsy and other seizure disorders, page 623.
84. Aarts WFM, Visser IH, Loonen, M.C.B, et al: (1998), Follow-up of 146 children with epilepsy after withdrawal of antiepileptic therapy. *Epilepsia*, 29: 244-250.
83. Abou-Khalil B, Andermann E., et al. (1993). Temporal lobe epilepsy after prolonged febrile convulsions excellent outcome after surgical treatment. *Epilepsia*, 34, 878-83.
84. Acharya J.N., Wyllie E, Laiders, H.O., et al (1997). Seizure symptomatology in infants with localization related epilepsy. *Neurology*, 48, 189-196.
85. Aicardi J, (1988), Clinical approach to the management of intractable epilepsy. *Developmental Medicine and Child Neurology*, 30: 429-440.
  - (1990a) Epilepsy in brain-injured children. *Developmental Medicine and Child Neurology*. 32:191-202.
  - (1990b), Neonatal myoclonic encephalopathy and early infantile epileptic encephalopathy.
    - In: Wasterlain, C.G., Verr. P. (Eds). *Neonatal Seizures*. New York: Raven Press, pp.41-49.
    - (1991), Neonatal seizures. In: Dam, M., Gram, L, (Eds.). *Comprehensive Epileptology*. New York: Raven Press, pp 99-112.
    - (1994a), *Epilepsy in Children*, 2<sup>nd</sup> Edn. New York: Raven Press.
    - (1994b), Syndromic classification in the management of childhood epilepsy. *Journal of Child Neurology*, 9, Suppl.2, 2S14-2S18).
    - (1994c), The place of neuronal migration abnormalities in children neurology. *Canadian Journal of Neurological Science*, 21, 185-193.
    - (1995), Typical absences in the first two years of life. In: Duncan, J., London: Churchill Communication Europe, pp.284-288.
    - (1996), Epileptic syndromes with onset in early childhood. In Wallace, S.J. (Ed.), *Epilepsy in Children*. London: Chapman & Hall, pp.247-274,
    - (1997), Epilepsy as a nonproxysmal disorder. *Acta Neuropediatrica* (in press).
86. Chevric, J.J. (1970 'Convulsive status epilepticus in infants and children.' *Epilepsia* 11, 187-197.
87. Shorvon, S. (1998) "Intractable epilepsy" In: Engel J., Pedley, TA) Eds). 'Epilepsy: A comprehensive Textbook Vol.2, New York: Lippincott-Raven, pp.1325.1331.
88. Taylor D.C. (1998), 'History and physical examination.' In Engel, J., Pedley, T.A. (Eds). *Epilepsy: A comprehensive Textbook Vol.I*, New York: Lippincott-Raven, pp. 805-810.



89. Sabril IS, Investigator and Peer Review Groups, et al. (1996). 'Vigabatrin
90. Caplan R. Epilepsy Syndromes. In: Coffey CE, Brumback RA (eds). Text Book of Pediatric Neuropsychiatry, 1st Edn. Washington DC, American Psychiatry Press, 1998, pp. 977-1010.
91. Engel J., Pedley TA, eds. Epilepsy: A comprehensive Textbook. New York, Lippincott-Raven, 1997.
92. Epilepsy. In: Swaiman KF, Ashwal S (eds), Pediatric Neurology: Principles and Practice, 3<sup>rd</sup> edn. St. Louis, Mosby, 1999, pp. 629-746.
93. Mankes JM, Sankar R. Paroxysmal disorder. In: Menkes JH (ed). Textbook of Child Neurology, 5<sup>th</sup> Edn. Baltimore, Williams and Wilkins, 1995, pp. 725-814.
94. Shinnar S, Amir N, Branski D. Childhood Seizures: Pediatric and Adolescent Medicine, Vol. 6, New York, S. Karger Publishing, 1995.
95. Berkovic, S.F., Howell, R.A., Hay, D.A. and Hoper J.L. Twin birth is not a risk factor for seizures. Neurology 1993; 43: 2515-2519.
96. Berkovic, S.F., Howell, R.A., Hay, D.A. and Hopper, J.L. Epilepsies in twins. In: Epileptic Seizures and Syndromes (Ed P. Wolf), Libbey, 1994: pp. 157-164.
97. Ottman, R. Genetic and developmental influences on susceptibility to epilepsy: evidence from twins. Paediatric and Perinatal Epidemiology 1992; 6: 265-272.
98. Churchill, J.A. The relationship of epilepsy to twin birth. Transactions of the American Microscopy Society 1958; 83: 118-120.
99. Sharma, K. Higher risk of epilepsy in twins. Indian Journal of Pediatrics 1986; 53: 515-519.
100. Commission of Classification and Terminology of the International League Against Epilepsy: Proposal for classification of epilepsies and epileptic syndromes. Epilepsia 1989; 30: 389-399.
101. Goulatia, R.K. Verma A., Misra, N.K. and Ahuja G.K. Disappearing CT lesions in epilepsy. Epilepsia 1987; 28: 523-527.
102. Bildhaiya, G.S. A study of twin births. Indian Pediatrics 1978; 15: 931-934.
103. Das Chaduhuri, A.B., Basu, S. and Chakraborty, S. Twinning rate in the Muslim population of West Bengal. Acta Genetica Medica Gemellologica Roma 1993; 42: 35-39.
104. Dutta, I. A study of twin births in Jabalpur city. Indian Journal of Pediatrics 1972; 39: 15-19.
105. Ghosh, S. and Ramajujacharyulu, T.K.T.S. Study of twin births in an urban community of Delhi. Indian Journal of Pediatrics 1979; 70: 70-77.
106. Joseph B, Hazards of multiple pregnancy. Journal of Obstetrics and Gynecology India 1964; 14: 118-127.
107. Junnarkar, A.R. and Nadkarni, M.G. Incidence of multiple births in an Indian rural community. Journal of Epidemiology and Community Health 1979; 33: 305-306.
108. Nylander, P.P.S.. The phenomenon of twinning. In: Obstetrical Epidemiology (Eds S.L. Barron and A.M. Thomson). London, Academic Press, 1983: pp. 143-165.
109. Park J.E. Observations of twin births in Gwalior City. Indian Journal of Pediatrics 1966; 33: 39-43.

110. Patel V.G. and Patel, B.C. A review of the cases of multiple pregnancy. *Journal of Obstetrics and Gynecology India* 1962; 12: 677-690.
111. Rao, P.S.S., Inbaraj, S.G. and Muthurathnam, S. Twinning rates in Tamil Nadu. *Journal of Epidemiology and Community Health* 1983; 37: 117-120.
112. Shah, S.B. and Patel, D.N. Twinning and structural defects. *Indian Paediatrics* 1984; 21:475-478.
113. Srivastav, J.R., Bhalla, M. and Bhalla, J.N. A study of twins. *Indian Paediatrics* 1977; 14: 121-127.
114. Trivedi, R.R. and Motashaw, ND. Outcome of the second of twins in 583 twin deliveries. *Journal of Obstetrics and Gynecology India* 1965; 15: 591-598.
115. Anderson VE, Rich SS, Hauser WA, Wilcox KJ. In: Anderson VE, Hauser WA, Leppik IE, Noebels JL, Rich SS, eds. *Genetic strategies in epilepsy research*. Amsterdam: Elsevier, 1991:89-103.
116. Anderson VE, Wilcox KJ, Rich SS, Leppik IE, Hauser W.A. Twin studies in epilepsy. In: Beck-Mannagetta G, Anderson VE, Doose H, Janz D, eds. *Genetics of the epilepsies*. Berlin: Springer-Verlag, 1989:145-55.
117. Leppert M, Anderson VE, Quattlebaum T, et al. Benign familial neonatal convulsions linked to genetic markers on chromosome 20. *Nature* 1989; 337: 647-8.
118. Malafosse A, Leboyer M, Dulac L., et al. Confirmation of linkage of benign familial neonatal convulsions to D20S20. *Hum Genet* 1992; 89:54-8.
119. Lewis TB, Leach RJ, Ward K, O'Connell P, Ryan SG. Genetic heterogeneity in benign familial neonatal convulsions; identification of a new locus on chromosome 8q. *Am J Hum Genet* 1993; 53: 670-5.
120. Tahvanainen E, et al. The gene for a novel recessively inherited human childhood epilepsy with progressive mental retardation maps to the distal short arm of chromosome 8. *Proc Natl Acad Sci USA* 1994; 91:7267-70.
121. Lehesjoki A-E, Koskiniemi M, Sistonen P, et al. Localization of a gene for progressive myoclonus epilepsy to chromosome 21q22. *Proc Natl Acad Sci USA* 1991; 88:3696-9.
122. Greenberg DA, Delgado-Escueta AV, Widelitz H, et al. Juvenile myoclonic epilepsy (JME) may be linked to the Bf and HLA loci in human chromosome 6. *Am J Genet* (1988; 31:185-92).
123. Weissebecker K, Durner M, Janz D, et al. Confirmation of linkage between juvenile myoclonic epilepsy locus and the HLA region of chromosome 6. *Am J Med Genet* 1991; 38:32-6.
124. Durner M, Sander T, Greenberg DA, Johnson K, Beck-Mannagetta G, Janz D. Localisation of idiopathic generalized epilepsy on chromosome 6p in families of juvenile myoclonic epilepsy patients. *Neurology* 1991; 41:1651-55.
125. Whitehouse W, Rees M, Curtis D, et al. Linkage analysis of idiopathic generalized epilepsy (IGE) and marker loci on chromosome 6p in families of patients with juvenile myoclonic epilepsy: no evidence for an epilepsy locus in the HLA region. *Am J Hum Genet* 1993; 55:652-62.
126. Ottman R, Risch N, Hauser W.A. et al. Localization of a gene for partial epilepsy to chromosome 10a. *Nature Genet* 1995; 10:56-60.



127. Phillips HA, Scheffer IE, Berkovic SF, et al. Localization of gene for autosomal dominant nocturnal frontal lobe epilepsy to chromosome 20q13.2. *Nature Genet* 1995;10:117-8.
128. McNamara JO. Identification of genetic defect of an epilepsy strategies for therapeutic advances. *Epilepsia* 1994; 35 (suppl 1): S51-7.
129. Roses AD, Paricak-Vance MA, Saunders AM et al. Complex genetic disease: Can genetic strategies in Alzheimer's disease and new genetic mechanisms be applied to epilepsy? *Epilepsia* 1994; 35 (suppl 1):S20-8.
130. Wolf P. Epilepsy with grand mal on awakening. In: Roger J., Dravet C, Bureau M, Dreifuss FE, Wolf P, eds. *Epileptic syndromes in infancy, childhood and adolescence*. London: John Libbey Eurotext, 1985:259-71.
131. Bhargava S, Tandon PN. Intracranial tuberculomas: a CT study. *J. Radiol* 1980; 53:935-45.
132. Rajshekhar V. Etiology and management of single small CT lesions in patients with seizure: understanding a controversy. *Acta Neurol Scan* 1991; 84:465-70.
133. Ahuja GK, Behari M, Goulatia RK, Jailkhani BL. Disappearing CT lesions in epilepsy; is tuberculosis or cysticercosis the cause? *J.Neurol Neurosurg Psychiatry* 1989; 52: 915-16.
134. Chandy MJ, Rajshekhar V, Ghosh S, et al. Single small enhancing CT lesions in Indian patients with epilepsy: considerations. *J.Neurol Neurosurg Psychiatry* 1991; 54: 702-5.
135. Ottman R, Lee JH, Risch N, Hauser WA, Susser M. Clinical indicators of genetic susceptibility of epilepsy. *Epilepsia* 1996; 37: 353-61.
136. Evans JH. Post-traumatic epilepsy. *Neurology* 1962; 12:665-74.
137. Caveness WF. Onset and cessation of fits following craniocerebral trauma. *J.Neurosurg* 1963; 10:570-82.
138. Caveness WF, Meirowsky AM, Rish BL, et al. The nature of posttraumatic epilepsy. *J.Neurosurg* 1979; 50:545-53.
139. Rimoin DL, Metrakos JK. La genétique des manifestations convulsives chez des familles hémiplegiques. In: Cauverier J International. *Second International Conference of Human Genetics. International Congress Series No.32*, London: Excerpta Medica 1961: Abstract 195.
140. Jain S, Mishra NK, Maheshwari MC. Disappearing CT lesions in epileptic patients – Indian experience. In: Sinha KK ed. *Progress in clinical neurosciences*. Ranchi: Neurological Society of India. 1990:25-32.

- Appendix

## Appendix-1

Performa used for the study of Epilepsy in children attending Railway central hospital New Delhi for degree of Doctor of Philosophy in Paediatric Medicine

Name.....Age/Sex.....reg.no.....

DATE OF BIRTH.....

Address.....

.....Pin

code.....

Tele.no.....Res.....Office.....

Occupation...Education level.....

Past history of febrile Seizures.....

If yes, age of onset...Total FC.....

Present Seizure type.....

Seizure frequency..... <1year      1-6 /year      >6/year

History of Todds Palsy.....

Treatment (name and dose per day).....

Seizure control      No Seizures      1-4 Sz      >then 4 Seizures

EEGdetails.....

If abnormal: ..... Generalised/focal/Focal with Generalised

CT/MRI ..... Normal/Abnormal

If Abnormal conclusion.....

Family history.....present/absent

Family pedigree.....

.....

If present Diagnostic Code.....

Final diagnosis and code...

Any other information-----

**Appendix-2**

**Seizure Classification to be used for study of Epilepsy in children in a subset of population for degree of Doctor of Philosophy in Paediatric Medicine from Bundelkhand university, Jhansi.**

**SEIZURE CLASSIFICATION**

- 1.1 Idiopathic- LRES**
- 1.2 Symptomatic- LRES, CT/MRI: Abnormal**
- 1.3 Cryptogenic- LRES, CT/MRI: Normal**

**Generalized Epileptic Syndromes (GES)**

- 2.1 Idiopathic age related syndromes**
- 2.2 Cryptogenic/Symptomatic**
  - West Syndrome/LG Syndrome**
  - With GTCS, MJ/ Abs. etc.**
- 2.3 symptomatic Epilepsies/ syndromes with GTCS as presenting feature**
- 3.0 Unclassified: Focal/ Generalized, Neonatal Sz, Others etc. Special Syndromes**
- 4.1 Febrile Convulsions (FC)**
- 4.2 Single Sz (SSZ)**
- 4.3 Situation related (SR) Sz associated with Alcohol / drugs / Ecampsia / Diabetes Ketoacidosis / Acute Metabolic events**
- 5.0 Seizure with Single Small Enhancing CT Lesion (SSEL)**
- 6.0 Others H/O Sz + but patients not seen dead relatives with H/o Sz.**

## RAILWAY NETWORK OF INDIA

